

Original Research Articles

Development of a Culturally Adaptable Intermediate Care Model For Older Adult Stroke Survivors

Temitope Farombi, MBBS, MSc, PhD¹, Hafiz T. A. Khan, PhD², Muili Lawal, PhD³

¹ Neurology, Chief Tony Anenih Geriatric Center, University College Hospital, Ibadan, Nigeria., ² College of Nursing, Midwifery, and Healthcare, University of West London, ³ College of Nursing, Midwifery and Healthcare, University of West London

Keywords: older adults, stroke, intervention, intermediate care

<https://doi.org/10.65104/001c.168525>

Journal of Brain Health

Vol. 1, Issue 2, 2026

Introduction

Intermediate care is vital for maximizing post-stroke functional recovery and maintaining a continuum of rehabilitation. However, robust evidence validating the development or practical deployment of home-based intermediate care models for older adults in Nigeria is currently absent.

Objective(s)

To develop and validate a home-based intermediate care model for improved functional outcomes and quality of life in older adult stroke survivors.

Methods

This study employed a two-round Delphi consensus process involving stakeholders in stroke care, including clinicians from geriatrics, neurology, nursing, physiotherapy, and medical social work, to develop and validate a home-based intermediate care model for family caregivers of stroke survivors. The model was developed through a multi-stage process. First, a systematic review was conducted to identify existing intermediate care models and their gaps. Subsequently, a qualitative study explored stakeholders' perspectives on the home-based care needs of older adult stroke survivors. Findings from the systematic review and qualitative study were integrated to develop a preliminary intermediate care model, which was subsequently refined and validated through the Delphi consensus process.

Results

The development and validation of the intermediate care model resulted in 10 domains of caregiving. These include stroke knowledge, seizure management, medication management, mobility and fall prevention, vital signs, service referral, reminiscence therapy, speech therapy, nutrition, and financial education.

Discussion

This model provides a structured framework to facilitate the transition of hospital to home care for older adult stroke survivors.

INTRODUCTION

Stroke is a global public health challenge that leads to disability, dementia, mortality, and carries significant economic impacts.^{1,2} Increases in stroke prevalence have been associated with an increase in the ageing population worldwide.^{3,4} Older adult stroke survivors, defined by WHO as those aged 60 years and older, face challenges including physical disabilities, cognitive and speech impairments, high rates of depression, and severe difficulty accessing consistent community post-stroke care and financial sup-

port.^{5,6} These challenges are worsened in Africa where substantial disparities and health inequality exist.^{7,8}

In Nigeria, stroke patients are often discharged without clearly communicated care plans for optimal functional recovery.⁹ Family caregivers of these patients often experience challenges in providing support and have poor knowledge and skills required to perform care activities.¹⁰ These systemic challenges highlight the urgent necessity for structured clinical guidelines to ensure adequate and appropriate care to reduce hospital readmissions. The intermediate care concept was introduced to improve the coordination of adequate care and safety during hospital

Plain Language summary

What is already known?

Stroke is a major cause of disability among older adults. Yet many survivors, especially in low-resource settings, lack access to structured post-discharge rehabilitation and caregiver support, leading to poor recovery outcomes.

What are the new findings/contributions?

Evidence from a systematic review and stakeholder interviews were used to design a practical home-based care model tailored to the needs of older stroke survivors and their caregivers. The model focused on caregiver education and support in areas such as stroke awareness, vital signs, medication use, mobility and fall prevention, nutrition, speech therapy, seizure management, reminiscence therapy, financial management, and service referral.

How might this impact clinical/care practice?

The model could improve continuity of care after hospital discharge by providing caregivers with the knowledge and support needed to manage stroke survivors at home. It could also contribute to better recovery outcomes and support the integration of home-based stroke care into routine clinical and community practice.

discharge.^{11,12} The development of intermediate care is intended to promote safe and early discharge out of acute care, foster independence, and enhance reintegration in community settings.^{13,14}

Basic intermediate care models include an integrated assessment and rehabilitation.¹⁵ Comparable to acute stroke intervention, intermediate care after stroke could improve longevity and quality of life.^{16,17} Evidence has shown that home-based intermediate care models, incorporated with training interventions for family caregivers, produce beneficial outcomes such as reduced risk of stroke recurrence and reduced mortality.^{18,19} Despite these potential benefits, there is little or no information on the development and implementation of intermediate care models for older adult stroke survivors in Nigeria.

OBJECTIVE(S)

This study aimed to develop and validate a home-based intermediate care model for improved functional outcomes and quality of life in older adult stroke survivors.

METHODS

STUDY DESIGN

A methodological study involving the development and expert validation of a culturally adaptable intermediate care model using a two-round Delphi consensus process.

DEVELOPMENT OF THE PRELIMINARY INTERMEDIATE CARE MODEL

The development of the intermediate care model followed a mixed method approach ([Figure 1](#)). In the first stage, a systematic review of existing intermediate care models was conducted and reported following the PICO framework²⁰ and the Preferred Reporting Items for Systematic Reviews

and Meta-Analyses (PRISMA) guideline.²¹ Eleven studies were included in the review (See Supplemental Materials), which revealed diverse interventions employed in the rehabilitation of older adult stroke survivors, encompassing exercise programs, speech therapy, video-based programs, reminiscence therapy, and caregiver-assisted therapy. The interventions demonstrated beneficial effects on functional outcomes and overall quality of life of participants.²² Stage two of the process included the use of an explorative design to obtain in-depth views from 18 family caregivers of stroke survivors who participated in focus group discussion (3 focus group discussions with 6 caregivers per group). Key informant interviews were also carried out with 8 medical experts, and 8 community gatekeepers ([Figure 2](#)). Thematic analysis was conducted using NVivo (version 12 pro) software and the coding technique used was inductive-dominant approach. Inclusion criteria for this stage were: family caregivers who have a dominant caring role in assisting older adult stroke survivors aged 60 years and older; medical experts involved in stroke care; and community gatekeepers who have had contact with at least one stroke survivor. This was conducted to understand the perceived needs of older adult stroke survivors. Insights from the study demonstrated that intermediate care should encompass physiotherapy service, minor medical treatment, nutritional service, speech therapy, mood disorder management, financial management, and medication assistance.²³ Following this stage, the preliminary intermediate care model ([Figure 2](#) and Supplemental Materials) was developed by integrating evidence from the both the systematic review and the qualitative study.

EXPERT VALIDATION

Selection and recruitment of experts: A multidisciplinary team of five medical professionals from a tertiary hospital were invited to participate in the review of the preliminary intermediate care model for accuracy and ease of use for




Figure 1. Process of Preliminary intermediate care model development




Figure 2. Preliminary intermediate care model.

older adult stroke survivors and their caregivers at home. The team comprised a consultant neurologist, a geriatric doctor, a geriatric nurse, a medical social worker, and a physiotherapist.

Validation tool/questionnaire: A 10-item questionnaire was developed to obtain experts' opinions regarding the adequacy, relevance, and completeness of each domain of the preliminary intermediate care model. Each item was rated using a five-point Likert scale ranging from strongly dis-

agree (1) to strongly agree (5). Open-ended comment boxes accompanied each item to enable experts to provide qualitative feedback and suggest modifications to improve the model (See Supplemental Materials).

Consensus criteria: Consensus was predefined as at least 100% of experts rating an item as agree or strongly agree.²⁴ Domains that did not achieve the consensus threshold or received substantial qualitative comments were revised before the subsequent Delphi round.

Validation procedure: A two-round Delphi consensus process was conducted to validate the preliminary model. Experts independently reviewed the model and completed a structured questionnaire during each round. Feedback obtained in the first round was incorporated into the revised model before redistribution for the second round.

DATA ANALYSIS

Quantitative data were analysed using descriptive statistics. The Item-Level Content Validity Index (I-CVI) was calculated as the proportion of experts who rated each item as either agree or strongly agree. The Scale-Level Content Validity Index based on the Average method (S-CVI/Ave) was calculated as the average of the I-CVI values across all items, while the Scale-Level Content Validity Index based on Universal Agreement (S-CVI/UA) was calculated as the proportion of items that achieved universal agreement among the experts.

ETHICAL CONSIDERATIONS

This work is part of a larger study with ethical approval from the University of Ibadan/University College Hospital (UI/UCH) Ethics Committee to ensure study compliance with the Helsinki's declaration. The UI/UCH Ethics Committee assigned number is UI/EC/22/0410.

RESULTS

A total of 5 medical experts participated in the first Delphi round, the Item-Level Content Validity Index (I-CVI) ranged from 0.80 to 1.00. Six of the ten domains achieved an I-CVI of 0.80, while the remaining four domains achieved an I-CVI of 1.00. The overall Scale-Level Content Validity Index/Average (S-CVI/Ave) was 0.88, indicating good overall content validity. However, the Scale-Level Content Validity Index/Universal Agreement (S-CVI/UA) was 0.40, reflecting that universal agreement was achieved for four of the ten domains only (Table 1).

Following the first round, the model was revised based on expert comments. Only one expert made comments, therefore, no formal qualitative analysis was performed. The comments were incorporated into the model before the second Delphi round. The principal recommendations were to strengthen the medication management domain by including a comprehensive list of the stroke survivor's medications to facilitate communication during hospital visits and consultations with other healthcare providers, ensuring that caregivers obtain and document daily vital signs measurements (*i.e.*, blood pressure, pulse, temperature, and respiratory rate) to track changes, promoting a healthy lifestyle, encouraging the children and family to be involved in assisting the patient financially.

The revised model was redistributed to the experts for the second Delphi round. In the second round, all ten domains achieved an I-CVI of 1.00, indicating 100% universal agreement. Similarly, both the S-CVI/Ave and S-CVI/UA

reached 1.00 (Table 2), demonstrating excellent content validity and complete consensus on the final model.

The final validated model (Table 3) was translated into an audio-visual resource (URL in Supplemental Materials) to complement the written intermediate care model (Supplemental Materials). This adaptation was implemented to optimize family caregivers' comprehension of the intervention guidelines and facilitate adherence to the guidelines within the home setting.

DISCUSSION

These findings indicate that the iterative Delphi process successfully improved the content validity and acceptability of the model. The development and validation of the intermediate care model identified 10 caregiving domains: stroke knowledge, seizure management, medication management, mobility and fall prevention, vital signs monitoring, service referral, reminiscence therapy, speech therapy, nutrition, and financial education. These domains are important because stroke survivors often experience deficits across multiple functional and quality-of-life areas, leaving caregivers feeling unprepared for their roles.²⁵ As stroke severity and recovery needs differ, personalized post-discharge care is essential.²⁶ Adequate caregiver education may therefore improve recovery outcomes and reduce caregiver burden.

The model was validated using the Delphi technique, which is widely applied in stroke intervention research.^{27, 28} Consensus was reached among a multidisciplinary panel consisting of a geriatric doctor, neurologist, geriatric nurse, medical social worker, and physiotherapist. The model was informed by evidence-based literature and stakeholder perspectives on intermediate stroke care, consistent with previous studies that developed intervention manuals from literature reviews.²⁹

Our findings are consistent with previous studies in Global South regions that developed intervention packages for stroke survivors and their caregivers. Similar interventions developed in India included education on stroke recovery trajectory, risk identification and management of low mood, task-specific rehabilitation activities, goal setting involving the patient and family caregiver, mobility training, and discharge planning.^{30,31} In comparison, our study developed a broader caregiver-focused model that emphasized stroke knowledge, medication management, and service referral.

The stroke knowledge domain in our intervention addressed stroke risk factors, symptoms, and prevention strategies, which aligns with earlier studies highlighting the importance of educating caregivers on stroke recovery and prevention. This is particularly important because inadequate caregiver knowledge and skills in post-stroke care may place both caregivers and stroke survivors at risk.^{32, 33} In addition, while previous interventions emphasized rehabilitation activities such as mobility and task-oriented training, our study further incorporated medication management, recognizing that many caregivers experience

Table 1. Outcomes of First Round

S/N	Item	Expert 1	Expert 2	Expert 3	Expert 4	Expert 5	Expert in agreement	I-CVI	UA
1	Stroke knowledge and prevention	1	1	0	1	1	4	0.8	0
2	Vital signs	1	1	0	1	1	4	0.8	0
3	Physiotherapy/ Mobility	1	1	0	1	1	4	0.8	0
4	Speech therapy	1	1	0	1	1	4	0.8	0
5	Reminiscence therapy	1	1	1	1	1	5	1	1
6	Financial management education	1	1	1	1	1	5	1	1
7	Nutritional service	1	1	0	1	1	4	0.8	0
8	Medication management	1	1	0	1	1	4	0.8	0
9	Seizure management	1	1	1	1	1	5	1	1
10	Service referral	1	1	1	1	1	5	1	1
							S-CVI/Ave	0.88	
	Proportion Relevance	1	1	0.4	1	1	S-CVI/UA		0.40
	Average proportion of items judged as relevance across the five experts						0.88		

I-CVI = Item-level Content Validity Index; UA = Universal Agreement; S-CVI/Ave = Scale-level Content Validity Index based on the Average; S-CVI/UA = Scale-level Content Validity Index based on Universal Agreement.

Table 2. Outcomes of Second Round

S/N	Item	Expert 1	Expert 2	Expert 3	Expert 4	Expert 5	Expert in agreement	I-CVI	UA
1	Stroke knowledge and prevention	1	1	1	1	1	5	1	1
2	Vital signs	1	1	1	1	1	5	1	1
3	Physiotherapy/ Mobility	1	1	1	1	1	5	1	1
4	Speech therapy	1	1	1	1	1	5	1	1
5	Reminiscence therapy	1	1	1	1	1	5	1	1
6	Financial management education	1	1	1	1	1	5	1	1
7	Nutritional service	1	1	1	1	1	5	1	1
8	Medication management	1	1	1	1	1	5	1	1
9	Seizure management	1	1	1	1	1	5	1	1
10	Service referral	1	1	1	1	1	5	1	1
							S-CVI/Ave	1	
	Proportion relevance	1	1	1	1	1	S-CVI/UA		1
	Average proportion of items judged as relevance across the five experts						1		

I-CVI = Item-level Content Validity Index; UA = Universal Agreement; S-CVI/Ave = Scale-level Content Validity Index based on the Average; S-CVI/UA = Scale-level Content Validity Index based on Universal Agreement.

Table 3. Domains and Content of the Final Intermediate Care Model

Intermediate Care Model developed for informal caregivers of older adult stroke patients	
Domain	Content
Stroke knowledge	- Definition of stroke - Symptoms of stroke - Types of stroke - Risk factors of stroke - Stroke prevention and management
Mobility and fall prevention	- Physiotherapy techniques
Vital signs	- Blood pressure measurement - Pulse rate - Body temperature - Respiration rate - Documentation of measurements - Visit to the healthcare provider
Speech therapy	- Visual and verbal cueing - Picture matching - Sentence completion task
Reminiscence therapy	- General guidelines - Picture matching - Story telling
Nutritional service	- Healthy foods for stroke patients - Unhealthy foods to avoid - Feeding techniques
Medication management	- General guidelines - Timely medication use - Use of medication discrepancy tool - Comprehensive list of medication
Seizure management	- Guidelines and first aid measures
Financial management	- Use of Health Insurance Scheme - Financial support by family members
Service referral	- Specialist care

challenges with the safe and timely administration of medications after discharge.³⁴

Another important distinction in our study was the inclusion of a service referral domain, which emphasized timely access to specialist care. This domain reinforced the importance of early treatment-seeking behavior among stroke survivors and caregivers when complications arise beyond the scope of home-based management.³⁵ While discharge planning was also highlighted in previous studies, our intervention extended this concept by explicitly guiding caregivers on when and where to seek specialized rehabilitation and supportive services. Overall, both our findings and previous studies underscore the importance of multi-dimensional caregiver education in improving continuity of care and supporting post-stroke recovery.

Although experts recommended additional clarification in some areas, consensus was reached to retain all domains in the model, suggesting their relevance to improving stroke outcomes. To enhance usability, a 25-minute audio-visual recording demonstrating the training content was developed. Previous studies have also explored technological approaches for caregiver interventions in stroke care.^{36, 37}

Task-shifting to caregivers in resource-constrained settings may reduce hospital admissions and improve outcomes for stroke survivors.³⁸ The proposed model offers a cost-effective approach for stroke care in Nigeria, where ac-

cess to rehabilitation infrastructure and skilled personnel remain limited. While the intervention has not yet been implemented, the development and validation process suggest it is conceptually appropriate for home-based stroke care, particularly in settings where family members provide most caregiving. However, effective implementation will require caregiver training, healthcare coordination, and access to rehabilitation resources.³⁹ Notably, caregiving responsibilities are often disproportionately borne by women, which may increase caregiver burden and reinforce existing inequities.^{40,41} Future research should therefore examine caregiver experiences and explore strategies for developing more equitable and supportive care models.

STRENGTHS AND LIMITATIONS

A major strength of this study is the design of the intermediate care model which is perhaps the first of its kind in Nigeria. Limitations of the study include the use of a small number of professionals and the exclusion of community stakeholders in the validation process. Additionally, the use of a single language (English language) in the audio-visual resource may hinder the effective use of the package among non-English-speaking family caregivers. Future adaptations and translations of the care model into other

languages will be critical for maximizing the accessibility of these resources.

CONCLUSIONS

In conclusion, this study successfully developed and validated a culturally adaptable intermediate care model for older adult stroke survivors. This structured framework is poised to significantly bolster home care support post-discharge, thereby ensuring continuity of care. Future investigations must rigorously assess the model's feasibility, acceptability, and ultimate effectiveness in improving clinical outcomes for stroke survivors. Ultimately, these findings provide critical evidence that can inform policy development concerning integrated stroke care pathways and discharge planning for vulnerable older adults.

.....

ACKNOWLEDGEMENTS

The authors sincerely acknowledge all healthcare professionals who contributed to the success of the Delphi consensus process and qualitative interviews, for their valuable

time, expertise, and contributions to the development and validation of the home-based intermediate care model. We also appreciate the family caregivers and stakeholders whose experiences informed the development of the model.

AUTHOR CONTRIBUTIONS

Conceptualization of study Design- Temitope Farombi, Muili Lawal and Hafiz Khan. Data Collection- Temitope Farombi. Data Analysis - Temitope Farombi, Muili Lawal and Hafiz Khan. Critical review - Muili Lawal and Hafiz Khan. First draft - Temitope Farombi. Final draft - Temitope Farombi, Muili Lawal and Hafiz Khan.

DISCLOSURES

The authors report no conflicts of interest.

FUNDING

This study received no external funding.

Submitted: May 18, 2026 IST. Accepted: August 17, 2026 IST.
Published: September 04, 2026 IST.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

REFERENCES

1. Akinyemi RO, Ovbiagele B, Adeniji OA, et al. Stroke in Africa: profile, progress, prospects and priorities. *Nat Rev Neurol*. 2021;17(10):634-656. doi:[10.1038/s41582-021-00542-4](https://doi.org/10.1038/s41582-021-00542-4)
2. López-Espuela F, Pedrera-Zamorano JD, Jiménez-Caballero PE, et al. Functional Status and Disability in Patients After Acute Stroke: A Longitudinal Study. *Am J Crit Care*. 2016;25(2):144-151. doi:[10.4037/ajcc2016215](https://doi.org/10.4037/ajcc2016215)
3. Katan M, Luft A. Global Burden of Stroke. *Semin Neurol*. 2018;38(02):208-211. doi:[10.1055/s-0038-1649503](https://doi.org/10.1055/s-0038-1649503)
4. Wang H, Naghavi M, Allen C, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*. 2016;388(10053):1459-1544. doi:[10.1016/S0140-6736\(16\)31012-1](https://doi.org/10.1016/S0140-6736(16)31012-1)
5. Rudnicka E, Napierała P, Podfigurna A, Męczekalski B, Smolarczyk R, Grymowicz M. The World Health Organization (WHO) approach to healthy ageing. *Maturitas*. 2020;139:6-11. doi:[10.1016/j.maturitas.2020.05.018](https://doi.org/10.1016/j.maturitas.2020.05.018)
6. Zeng S, Wu M, Xu L, et al. Challenges in Accessing Community-Based Rehabilitation and Long-Term Care for Older Adult Stroke Survivors and Their Caregivers: A Qualitative Study. *J Multidiscip Healthc*. 2024;17:4829-4838. doi:[10.2147/JMDH.S476993](https://doi.org/10.2147/JMDH.S476993)
7. Aderinto N, Olatunji G, Kokori E, et al. Disparities in post-stroke rehabilitation: an African perspective. *Discov Public Health*. 2025;22(1):218. doi:[10.1186/s12982-025-00622-6](https://doi.org/10.1186/s12982-025-00622-6)
8. Feigin VL, Abajobir AA, Abate KH, et al. Global, regional, and national burden of neurological disorders during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Neurol*. 2017;16(11):877-897. doi:[10.1016/S1474-4422\(17\)30299-5](https://doi.org/10.1016/S1474-4422(17)30299-5)
9. Soeker MS, Olaoye OA. Exploring the experiences of rehabilitated stroke survivors and stakeholders with regard to returning to work in South- West Nigeria. *Work*. 2017;57(4):595-609. doi:[10.3233/WOR-172590](https://doi.org/10.3233/WOR-172590)
10. Bierhals CCBK, Santos NOD, Fengler FL, Raubustt KD, Forbes DA, Paskulin LMG. Needs of family caregivers in home care for older adults. *Rev Lat Am Enfermagem*. 2017;25(0). doi:[10.1590/1518-8345.1511.2870](https://doi.org/10.1590/1518-8345.1511.2870)
11. Chen L, Sit JWH, Shen X. Quasi-experimental evaluation of a home care model for patients with stroke in China. *Disabil Rehabil*. 2016;38(23):2271-2276. doi:[10.3109/09638288.2015.1123305](https://doi.org/10.3109/09638288.2015.1123305)
12. Dahl U, Steinsbekk A, Johnsen R. Effectiveness of an intermediate care hospital on readmissions, mortality, activities of daily living and use of health care services among hospitalized adults aged 60 years and older—a controlled observational study. *BMC Health Serv Res*. 2015;15(1):351. doi:[10.1186/s12913-015-1022-x](https://doi.org/10.1186/s12913-015-1022-x)
13. Lo YP, Wang MC, Chen YH, Chiang SL, Lin CH. Effectiveness of a Post-Acute-Care Rehabilitation Program in Patients with Stroke: A Retrospective Cohort Study. *Life*. 2025;15(8):1216. doi:[10.3390/life15081216](https://doi.org/10.3390/life15081216)
14. Sehatkarlangrodi S, Zargoush M, Longo CJ, Ghazalbash S. Shaping older adults' care policy: a scoping review of key determinants in post-acute and community reintegration transitions. *BMC Health Serv Res*. 2025;25(1):1236. doi:[10.1186/s12913-025-13433-x](https://doi.org/10.1186/s12913-025-13433-x)
15. Sezgin D, O'Caoimh R, O'Donovan MR, et al. Defining the characteristics of intermediate care models including transitional care: an international Delphi study. *Aging Clin Exp Res*. 2020;32(11):2399-2410. doi:[10.1007/s40520-020-01579-z](https://doi.org/10.1007/s40520-020-01579-z)
16. Duncan PW, Bushnell C, Sissine M, et al. Comprehensive Stroke Care and Outcomes: Time for a Paradigm Shift. *Stroke*. 2021;52(1):385-393. doi:[10.1161/STROKEAHA.120.029678](https://doi.org/10.1161/STROKEAHA.120.029678)
17. Tan SYJ, Lee SC, Rusli KDB, et al. Effectiveness of Multidisciplinary Transitional Care Interventions on Functional Status, Quality of Life and Readmission Rates in Stroke Patients: A Systematic Review and Meta-Analysis. *J Clin Nurs*. 2026;35(1):85-98. doi:[10.1111/jocn.70000](https://doi.org/10.1111/jocn.70000)
18. Alim M, Lindley R, Felix C, et al. Family-led rehabilitation after stroke in India: the ATTEND trial, study protocol for a randomized controlled trial. *Trials*. 2016;17(1):13. doi:[10.1186/s13063-015-1129-8](https://doi.org/10.1186/s13063-015-1129-8)

19. Wang Y, Yang F, Shi H, Yang C, Hu H. What Type of Transitional Care Effectively Reduced Mortality and Improved ADL of Stroke Patients? A Meta-Analysis. *Int J Environ Res Public Health*. 2017;14(5):510. doi:[10.3390/ijerph14050510](https://doi.org/10.3390/ijerph14050510)
20. Cumpston MS, McKenzie JE, Thomas J, Brennan SE. The use of "PICO for synthesis" and methods for synthesis without meta-analysis: protocol for a survey of current practice in systematic reviews of health interventions. *F1000Research*. 2021;9:678. doi:[10.12688/f1000research.24469.2](https://doi.org/10.12688/f1000research.24469.2)
21. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. Published online March 29, 2021:n71. doi:[10.1136/bmj.n71](https://doi.org/10.1136/bmj.n71)
22. Farombi TH, Khan HT, Lawal M, Akinyemi R. Impact of intermediate home-based care on functional health of older adults with stroke in low-income and middle-income countries: A systematic review. *J Alzheimers Dis Rep*. 2025;9:25424823251318227. doi:[10.1177/25424823251318227](https://doi.org/10.1177/25424823251318227)
23. Farombi TH, Khan HTA, Lawal M. Exploring the Experiences of Family Caregivers in the Management of Stroke Among the Older Adults in Nigeria: A Qualitative Study. *J Popul Ageing*. 2024;17(4):679-691. doi:[10.1007/s12062-024-09454-9](https://doi.org/10.1007/s12062-024-09454-9)
24. Polit DF, Beck CT, Owen SV. Is the CVI an acceptable indicator of content validity? Appraisal and recommendations. *Res Nurs Health*. 2007;30(4):459-467. doi:[10.1002/nur.20199](https://doi.org/10.1002/nur.20199)
25. Lin S, Wang C, Wang Q, et al. The experience of stroke survivors and caregivers during hospital-to-home transitional care: A qualitative longitudinal study. *Int J Nurs Stud*. 2022;130:104213. doi:[10.1016/j.ijnurstu.2022.104213](https://doi.org/10.1016/j.ijnurstu.2022.104213)
26. Fisher RJ, Walker MF, Golton I, Jenkinson D. The implementation of evidence-based rehabilitation services for stroke survivors living in the community: the results of a Delphi consensus process. *Clin Rehabil*. 2013;27(8):741-749. doi:[10.1177/0269215512473312](https://doi.org/10.1177/0269215512473312)
27. Andrades-González I, Molina-Mula J. Validation of Content for an App for Caregivers of Stroke Patients through the Delphi Method. *Int J Environ Res Public Health*. 2022;19(12):7523. doi:[10.3390/ijerph19127523](https://doi.org/10.3390/ijerph19127523)
28. Santos NOD, Predebon ML, Bierhals CCBK, Day CB, Machado DDO, Paskulin LMG. Development and validation a nursing care protocol with educational interventions for family caregivers of elderly people after stroke. *Rev Bras Enferm*. 2020;73(suppl 3):e20180894. doi:[10.1590/0034-7167-2018-0894](https://doi.org/10.1590/0034-7167-2018-0894)
29. Fuhrmann AC, Bierhals CCBK, Santos NOD, Machado DDO, Cordova FP, Paskulin LMG. CONSTRUCTION AND VALIDATION OF AN EDUCATIONAL MANUAL FOR FAMILY CAREGIVERS OF OLDER ADULTS AFTER A STROKE. *Texto Contexto - Enferm*. 2021;30:e20190208. doi:[10.1590/1980-265x-tce-2019-0208](https://doi.org/10.1590/1980-265x-tce-2019-0208)
30. Lindley RI, Anderson CS, Billot L, et al. Family-led rehabilitation after stroke in India (ATTEND): a randomised controlled trial. *The Lancet*. 2017;390(10094):588-599. doi:[10.1016/S0140-6736\(17\)31447-2](https://doi.org/10.1016/S0140-6736(17)31447-2)
31. Pandian JD, Felix C, Kaur P, et al. Family-Led Rehabilitation after Stroke in India: The ATTEND Pilot Study. *Int J Stroke*. 2015;10(4):609-614. doi:[10.1111/ijss.12475](https://doi.org/10.1111/ijss.12475)
32. Parappilly BP, Mortenson WB, Field TS, Eng JJ. Exploring perceptions of stroke survivors and caregivers about secondary prevention: a longitudinal qualitative study. *Disabil Rehabil*. 2020;42(14):2020-2026. doi:[10.1080/09638288.2018.1544296](https://doi.org/10.1080/09638288.2018.1544296)
33. Rumiati R, Kariasa IM, Waluyo A. The Effectiveness of Post-Stroke Patient Care Education Intervention in Stroke Caregivers: A Literature Review. *IJNP Indones J Nurs Pract*. 2021;5(2). doi:[10.18196/ijnp.v5i2.11437](https://doi.org/10.18196/ijnp.v5i2.11437)
34. Gibson J, Coupe J, Watkins C. Medication adherence early after stroke: using the Perceptions and Practicalities Framework to explore stroke survivors', informal carers' and nurses' experiences of barriers and solutions. *J Res Nurs*. 2021;26(6):499-514. doi:[10.1177/1744987121993505](https://doi.org/10.1177/1744987121993505)
35. Whitehead S, Baalbergen E. Post-stroke rehabilitation. *S Afr Med J*. 2019;109(2):81. doi:[10.7196/SAMJ.2019.v109i2.00011](https://doi.org/10.7196/SAMJ.2019.v109i2.00011)
36. Canto DFD, Costa FMD, Woiciechoski LR, Cogo ALP, Paskulin LMG. Development of an online course for caregivers of older stroke patients. *Rev Gaúcha Enferm*. 2023;44:e20230040. doi:[10.1590/1983-1447.2023.20230040.en](https://doi.org/10.1590/1983-1447.2023.20230040.en)
37. Caunca MR, Simonetto M, Hartley G, Wright CB, Czaja SJ. Design and Usability Testing of the Stroke Caregiver Support System: A Mobile-Friendly Website to Reduce Stroke Caregiver Burden. *Rehabil Nurs*. 2020;45(3):166-177. doi:[10.1097/RNJ.0000000000000196](https://doi.org/10.1097/RNJ.0000000000000196)
38. Nkoke C. Post-stroke care: an alternative model to reduce stroke related morbidity in sub-Saharan Africa. *Annals of Translational Medicine*. 2015;3(16):238. doi:[10.3978/j.issn.2305-5839.2015.09.09](https://doi.org/10.3978/j.issn.2305-5839.2015.09.09)

39. Boettger P, Juenemann M, Buerke M, Agarwal A, Omar OA. Global Inequalities in Stroke Rehabilitation: System Capacity, Workforce, and Access in Low- and Middle-Income Countries. *Curr Phys Med Rehabil Rep*. 2026;14(1):12. doi:[10.1007/s40141-026-00535-8](https://doi.org/10.1007/s40141-026-00535-8)
40. Chica-Pérez A, Correa-Casado M, Fernández-Sola C, Granero-Molina J, Cortés-Rodríguez AE, Hernández-Padilla JM. The Family Caregiver Role From the Perspective of Older Women Experiencing Poverty in a High-Income Country: A Qualitative Study. *J Adv Nurs*. 2026;82(4):3317-3330. doi:[10.1111/jan.17043](https://doi.org/10.1111/jan.17043)

41. Ong APR, De Jesus LL, Linog S, et al. Burden of informal caregiving for stroke survivors in the Philippines: a cross-sectional survey of sociodemographic and caregiving factors. *Disabil Rehabil*. Published online April 15, 2026:1-18. doi:[10.1080/09638288.2026.2653953](https://doi.org/10.1080/09638288.2026.2653953)

SUPPLEMENTARY MATERIALS

Supplemental Materials: Development of a Culturally Adaptable Intermediate Care Model for Older Adult Stroke Survivors

Supplemental Materials:

Audiovisual link

Preliminary intermediate care model

Final intermediate care model

Validation tool for Delphi process

Table of Studies included in the Systematic Review

Download: <https://www.jbrainhealth.org/article/168525-development-of-a-culturally-adaptable-intermediate-care-model-for-older-adult-stroke-survivors/attachment/361928.pdf>
